

Total $[1,2]$ -domination in graphs*

Xuezheng Lv

Department of Mathematics, Renmin University of China,
Beijing 100872, P.R. China

Baoyindureng Wu [†]

College of Mathematics and System Sciences, Xinjiang University
Urumqi, Xinjiang 830046, P.R. China

Abstract: A subset $S \subseteq V$ in a graph $G = (V, E)$ is a total $[1, 2]$ -set if, for every vertex $v \in V$, $1 \leq |N(v) \cap S| \leq 2$. The minimum cardinality of a total $[1, 2]$ -set of G is called the total $[1, 2]$ -domination number, denoted by $\gamma_{t[1,2]}(G)$.

We establish two sharp upper bounds on the total $[1, 2]$ -domination number of a graph G in terms of its order and minimum degree, and characterize the corresponding extremal graphs achieving these bounds. Moreover, we give some sufficient conditions for a graph without total $[1, 2]$ -set and for a graph with the same total $[1, 2]$ -domination number, $[1, 2]$ -domination number and domination number.

Keywords: total $[1, 2]$ -set; total $[1, 2]$ -domination number; $[1, 2]$ -set

1 Introduction

We consider undirected finite simple graphs only, and refer to [13] for undefined notations and terminologies. Let $G = (V(G), E(G))$ be a graph of order $n = |V(G)|$ and size $m = |E(G)|$. For a vertex $v \in V(G)$, the *neighborhood* $N_G(v)$ of v is the set of vertices adjacent to v in G , and the *closed neighborhood* $N_G[v]$ of v is $N_G(v) \cup \{v\}$; the *degree* $d_G(v)$ of v is the number of edges incident with v in G . Since G is simple, $d_G(v) = |N_G(v)|$. If there is no confusion, we simply write $N(v)$, $N[v]$ and $d(v)$ instead of $N_G(v)$, $N_G[v]$ and $d_G(v)$. The minimum degree and maximum degree of a vertex in

*This work was supported by National Natural Science Foundation of China (No. 11001269, No. 11161046)

[†]Corresponding author.

E-mail address: xzlv@ruc.edu.cn (X. Lv); wubaoyin@hotmail.com (B. Wu)

a graph G are denoted by $\delta(G)$ and $\Delta(G)$, respectively. A vertex of degree 1 is called a *leaf* and the vertex adjacent to a leaf is called a *support vertex*.

Let $S \subseteq V(G)$ and $v \in S$. The open and closed neighborhood of S are $N(S) = \cup_{v \in S} N(v)$ and $N[S] = \cup_{v \in S} N[v]$, respectively. The *S -private neighborhood* of v , denoted by $pri_G(v, S)$ or simply $pri(v, S)$, consists of all vertices in $N_G[v]$ but not in $N_G[S \setminus \{v\}]$; that is, $pri(v, S) = N_G[v] \setminus N_G[S \setminus \{v\}]$. Thus, if $u \in pri(v, S)$, then $N[u] \cap S = \{v\}$. For $S' \subseteq S$, denote $pri(S', S) = \bigcup_{u \in S'} pri(u, S)$. The subgraph obtained by deleting all vertices in S is denoted by $G - S$. The symbol $G[S]$ denotes the subgraph of G induced by S . So, $G - S$ is the subgraph of G induced by $V(G) \setminus S$.

Let X and Y be two sets of vertices of a graph G . We denote by $E[X, Y]$ the set of edges of G with one end in X and the other end in Y .

A set $S \subseteq V(G)$ is called a *dominating set* of a graph G if $N[S] = V(G)$, that is, $N(v) \cap S \neq \emptyset$ for every vertex $v \in V(G) \setminus S$. The minimum cardinality of a dominating set in a graph G is called the *domination number* of G , and is denoted by $\gamma(G)$. A dominating set S of G is called a *total dominating set* if for every vertex $v \in V(G)$, $N(v) \cap S \neq \emptyset$. The *total domination number* of G , denoted by $\gamma_t(G)$, is the minimum cardinality of a total dominating set of G . An independent set of G is a set of mutually independent vertices.

The notion of $[1, 2]$ -domination was investigated by Dejter [6] and more recently, by Chellali et al. [4]. A dominating set S of G is called a *$[1, 2]$ -set* if $1 \leq |N(v) \cap S| \leq 2$ holds for every vertex $v \in V(G) \setminus S$, that is, S is a dominating set and every vertex in $V \setminus S$ is adjacent to no more than two vertices of S . The *$[1, 2]$ -domination number* of G , denoted by $\gamma_{[1,2]}(G)$, is the minimum cardinality of a $[1, 2]$ -set of G . Since the vertex set $V(G)$ itself is a $[1, 2]$ -set of a graph G , any graph has a $[1, 2]$ -set. There exists an infinite family of graphs G whose $[1, 2]$ -domination number are equal to their orders [4, 14]. So, one of the fundamental problem on $[1, 2]$ -domination of graphs is that for which graphs G of order n is $\gamma_{[1,2]}(G) = n$? The problem, whether there is a simple, polynomial test for deciding if $\gamma_{[1,2]}(G) < n$, is still open. A number of open problems in [4] regarding $[1, 2]$ -domination were solved by Yang and Wu [14].

An analogue notion of $[1, 2]$ -domination for the total domination of a graph was also introduced in [4]. A total dominating set S of G is called a *total $[1, 2]$ -set* if for every vertex $v \in V(G)$, $1 \leq |N(v) \cap S| \leq 2$. The *total $[1, 2]$ -domination number* of G , denoted by $\gamma_{t[1,2]}(G)$, is the minimum cardinality of a total $[1, 2]$ -set of G . However, we will see that there exist an infinite number of graphs with no total $[1, 2]$ -set, even among trees. For convenience, if there is no total $[1, 2]$ -set in a graph G , we denote $\gamma_{t[1,2]}(G) = +\infty$. By

this convention, it is trivial to see that for any graph G ,

$$\gamma(G) \leq \gamma_t(G) \leq \gamma_{t[1,2]}(G) \text{ and } \gamma(G) \leq \gamma_{[1,2]}(G) \leq \gamma_{t[1,2]}(G).$$

We call $S \subseteq V(G)$ a $\gamma(G)$ -set, if S is a dominating set with $|S| = \gamma(G)$. Similarly, $\gamma_{[1,2]}(G)$ -set, $\gamma_t(G)$ -set, $\gamma_{t[1,2]}(G)$ -set can be defined. One of the remaining problems in [4] was stated as follows.

Question ([4]). What can you say about total $[1, 2]$ -set, and the corresponding total $[1, 2]$ -domination number $\gamma_{t[1,2]}(G)$?

In this paper, we give two sharp upper bounds on the total $[1, 2]$ -domination number of a graph in terms of its order and an additional condition that the minimum degree at least 1 or at least 2. Moreover, we give some sufficient conditions for a graph without total $[1, 2]$ -set and for a graph with the same total $[1, 2]$ -domination number, $[1, 2]$ -domination number and domination number.

2 Upper bounds for $\gamma_{t[1,2]}(G)$

As usual, the path, cycle, and complete graph of order n are denoted by P_n , C_n and K_n , respectively. The k -corona $G \circ P_k$ is the graph obtained from G by attaching a pendant path of length $k - 1$ to each vertex $v \in V(G)$; the double k -corona $G \circ 2P_k$ is the graph obtained from G by attaching two pendant paths of length k to each vertex $v \in V(G)$, see Fig. 1 for an illustration.




Fig. 1: 2-corona and double 2-corona

We start with the following lemma.

Lemma 2.1. *Let $n \geq 3$ and $n_i \geq 1$ be integers for $1 \leq i \leq 2$. Then*

$$(1) \quad \gamma_{t[1,2]}(K_n) = \gamma_{t[1,2]}(K_{n_1, n_2}) = 2;$$

(2)

$$\gamma_{t[1,2]}(P_n) = \gamma_{t[1,2]}(C_n) = \begin{cases} \frac{n+1}{2}, & n \equiv 1 \pmod{2} \\ \frac{n}{2} + 1, & n \equiv 2 \pmod{4} \\ \frac{n}{2}, & n \equiv 0 \pmod{4}. \end{cases}$$

(3) if $G = H \circ 2P_2$ for a connected graph H of order at least two, then $\gamma_{t[1,2]}(G) = \frac{4n}{5}$, where n is the order of G .

Proof. We leave (1) and (2) to the readers for an exercise. To show (3), let H be a connected graph of order $k \geq 2$, with $V(H) = \{u_1, \dots, u_k\}$. Let $V(G) = V(H) \cup X \cup Y \cup X' \cup Y'$, where $X = \{x_1, \dots, x_k\}$, $Y = \{y_1, \dots, y_k\}$, $X' = \{x'_1, \dots, x'_k\}$, $Y' = \{y'_1, \dots, y'_k\}$, and $E(G) = E(H) \cup \{u_i x_i \mid 1 \leq i \leq k\} \cup \{u_i y_i \mid 1 \leq i \leq k\} \cup \{x_i x'_i \mid 1 \leq i \leq k\} \cup \{y_i y'_i \mid 1 \leq i \leq k\}$. Obviously, $X \cup Y \cup X' \cup Y'$ is a total $[1, 2]$ -set of G , so $\gamma_{t[1,2]}(G) \leq \frac{4n}{5}$. Take a $\gamma_{t[1,2]}$ -set S of G . Since S is a total dominating set of G , $X \cup Y \subseteq S$. On the other hand, $V(H) \cap S = \emptyset$. If it is not true, without loss of generality, we may assume that $u_1 \in S$ and u_2 is a neighbor of u_1 in H . But then, $\{x_2, y_2, u_1\} \subseteq N_G(u_2) \cap S$, contradicting that S is a total $[1, 2]$ -set of G . Combining the above two facts with the other two facts that $X \cup Y$ is an independent set of G and $\delta(G[S]) \geq 1$, it follows that $S = X \cup Y \cup X' \cup Y'$, and thus $\gamma_{t[1,2]}(G) = \frac{4n}{5}$. $\square$

By Lemma 2.1, $\gamma_{t[1,2]}(G) \geq 2$ for any connected graph G of order $n \geq 2$, and there are infinite family of graphs G with $\gamma_{t[1,2]}(G) = 2$. Indeed, for a graph G , $\gamma_{t[1,2]}(G) = 2$ if and only if G has a dominating set, which consists of a pair of adjacent vertices in G . Cockayne et al. [5] proved that $\gamma_t(G) \leq \frac{2n}{3}$ for a connected graph of order $n \geq 3$. Later, Brigham et al. [3] characterized those graphs achieving this bound.

Theorem 2.2. ([3, 5]) *Let G be a connected graph of order $n \geq 3$. Then $\gamma_t(G) \leq \frac{2n}{3}$ and the equality holds if and only if G is C_3 , C_6 or $H \circ P_2$ for some connected graph H .*

Note that $\gamma_{[1,2]}(G) \leq n$ is a trivial upper bound for a graph G of order n , and the equality holds for an infinite many values of n , see [4, 14]. In the following theorem, we establish a sharp upper bound for the total $[1, 2]$ -domination number of a connected graph in terms of its order and characterize all graphs achieving the bound.

Theorem 2.3. *Let G be a connected graph of order $n \geq 5$. If $\gamma_{t[1,2]}(G) < +\infty$, then*

$$\gamma_{t[1,2]}(G) \leq \frac{4n}{5},$$

with equality if and only if $G = H \circ 2P_2$ for some connected graph H of order at least two.

Proof. Let S be a $\gamma_{t[1,2]}(G)$ -set of G . Since $1 \leq \delta(G[S]) \leq \Delta(G[S]) \leq 2$, each component of $G[S]$ is a path or a cycle. Divide S into four subsets:

$$S_1 = \{u \in S \mid u \text{ lies in a component of } G[S] \text{ isomorphic to a cycle}\},$$

$$S_2 = \{u \in S \mid u \text{ lies in a component of } G[S] \text{ isomorphic to } K_2\},$$

$$S_3 = \{u \in S \mid u \text{ lies in a component of } G[S] \text{ isomorphic to } P_3\},$$

$$S_4 = \{u \in S \mid u \text{ lies in a component of } G[S] \text{ isomorphic to a path of order at least 4}\}.$$

Clearly, $|S| = \sum_{i=1}^4 |S_i|$. Recall that for a vertex $u \in S$, $\text{pri}(u, S) = \{v \in V(G) \setminus S \mid N(v) \cap S = \{u\}\}$ and $\text{pri}(S_i, S) = \cup_{u \in S_i} \text{pri}(u, S)$.

Claim 1. (1) $|\text{pri}(S_1, S)| \geq |S_1|$, (2) $|\text{pri}(S_3, S)| \geq \frac{2|S_3|}{3}$, (3) $|\text{pri}(S_4, S)| \geq |S_4| - 2\omega_4$, where ω_4 denotes the number of components in $G[S]$, which is isomorphic to a path of order at least 4.

Proof of Claim 1. Let $u \in S$ be a vertex. If $u \in S_1$ and $\text{pri}(u) = \emptyset$, then $S \setminus \{u\}$ will be a total $[1, 2]$ -set of G , contradicting the choice of S . Therefore

$$|\text{pri}(S_1, S)| = \left| \bigcup_{u \in S_1} \text{pri}(u, S) \right| = \sum_{u \in S_1} |\text{pri}(u, S)| \geq |S_1|.$$

This proves (1).

Now assume that $u \in S_3$ and $d_{G[S_3]}(u) = 1$. If $\text{pri}(u) = \emptyset$, $S \setminus \{u\}$ is still a total $[1, 2]$ -set of G , contradicting the choice of S . Therefore

$$|\text{pri}(S_3, S)| = \left| \bigcup_{u \in S_3} \text{pri}(u, S) \right| \geq \sum_{u \in S_3, d_{G[S_3]}(u)=1} |\text{pri}(u, S)| \geq \frac{2|S_3|}{3}.$$

This proves (2).

To show (3), let $P = u_1 u_2 \cdots u_k$ be a component of $G[S]$, which is isomorphic to a path of order k , where $k \geq 4$. By an argument similar to the above, we have $\text{pri}(u_i, S) \neq \emptyset$ for any $i \in \{1, 2, \dots, k\} \setminus \{2, k-1\}$, and thus $|\text{pri}(S_4, S)| \geq |S_4| - 2\omega_4$. $\square$

Claim 2. $|U| \geq \frac{|S_2|}{4}$ where $U = V(G) \setminus [S \cup \text{pri}(S_1, S) \cup \text{pri}(S_3, S) \cup \text{pri}(S_4, S)]$.

Proof of Claim 2. Since G is connected, every component of $G[S]$, which is isomorphic to K_2 , has at least one neighbor in U . Let $W = \cup_{u \in S_2} N(u) \cap U$. Since every vertex of W is adjacent to at most two vertices of S in G ,

$$2|W| \geq |E[S, W]| \geq |E[S_2, W]| \geq \frac{|S_2|}{2},$$

and thus $|U| \geq |W| \geq \frac{|S_2|}{4}$. □

Since $\omega_4 \leq \frac{|S_4|}{4}$, we have

$$\begin{aligned}
n &= |S| + |pri(S_1, S)| + |pri(S_3, S)| + |pri(S_4, S)| + |U| \\
&\geq |S| + |S_1| + \frac{2|S_3|}{3} + |S_4| - 2\omega_4 + \frac{|S_2|}{4} \\
&\geq \frac{5|S|}{4} + \frac{3|S_1|}{4} + \frac{5|S_3|}{12} + \frac{3|S_4|}{4} - \frac{|S_4|}{2} \\
&\geq \frac{5|S|}{4} + \frac{3|S_1|}{4} + \frac{5|S_3|}{12} + \frac{|S_4|}{4} \\
&\geq \frac{5|S|}{4}.
\end{aligned}$$

So, $\gamma_{t[1,2]}(G) \leq \frac{4n}{5}$.

If the equality holds, we have $|S_1| = |S_3| = |S_4| = 0$, $|V(G) \setminus S| = \frac{|S|}{4}$, every component isomorphic to K_2 has exactly one neighbor in $V(G) \setminus S$, and each vertex of $V(G) \setminus S$ has exactly two neighbors in S . And if $\frac{|S|}{4} = 1$, then $G = P_5$, while $\gamma_{t[1,2]}(P_5) = 3$. So, $G = H \circ 2P_2$ for a connected graph H of order at least 2. By (3) of Lemma 2.1, the converse is also true. □

Theorem 2.3 tells us that there does not exist a graph G of order $n \geq 5$ with $\gamma_{t[1,2]}(G) = k$ for any $\lfloor \frac{4n}{5} \rfloor < k \leq n$. However, the situation become different when $k \leq \lfloor \frac{4n}{5} \rfloor$. To see this let us construct a class of the following graphs. We start from a complete graph K_{n-k} with $V(K_{n-k}) = \{v_1, v_2, \dots, v_{n-k}\}$ and denote $r = \lfloor \frac{k}{4} \rfloor$. For $10 \leq k \leq \lfloor \frac{4n}{5} \rfloor$ and $k \equiv 0 \pmod{2}$, we construct a graph $F_{n,k}$ of order n with $\gamma_{t[1,2]}(F_{n,k}) = k$ as follows: $F_{n,k}$ has

$$V(F_{n,k}) = V(H) \cup \{w_1, \dots, w_{\frac{k}{2}}\} \cup \{w'_1, \dots, w'_{\frac{k}{2}}\}$$

and $E(F_{n,k}) = E(H) \cup \{w_i w'_i \mid 1 \leq i \leq \frac{k}{2}\} \cup E'$, where

$$E' = \{v_i w_{2i-1}, v_i w_{2i} \mid 1 \leq i \leq r-1\} \cup \{v_r w_{2r-1}\} \cup \{v_j w_{2r}, v_j w'_{2r} \mid r+1 \leq j \leq n-k\}$$

if $k \equiv 0 \pmod{4}$; and

$$E' = \{v_i w_{2i-1}, v_i w_{2i} \mid 1 \leq i \leq r\} \cup \{v_j w_{2r+1} v_j w'_{2r+1} \mid r+1 \leq j \leq n-k\}$$

if $k \equiv 2 \pmod{4}$. For $10 \leq k < \lfloor \frac{4n}{5} \rfloor$ and $k \equiv 1 \pmod{2}$, we construct a graph $F_{n,k}$ of order n with $\gamma_{t[1,2]}(F_{n,k}) = k$ as follows. Note that now $n - k - r \geq 2$. $F_{n,k}$ has

$$V(F_{n,k}) = V(H) \cup \{w_1, \dots, w_{\frac{k-3}{2}}\} \cup \{w'_1, \dots, w'_{\frac{k-3}{2}}\} \cup \{w, w', w''\}$$

and $E(F_{n,k}) = E(H) \cup \{w_i w'_i \mid 1 \leq i \leq \frac{k-3}{2}\} \cup \{w w', w' w''\} \cup E'$, where

$$E' = \{v_i w_{2i-1}, v_i w_{2i} \mid 1 \leq i \leq r-1\} \cup \{v_r w_{2r-1}, v_{r+1} w, v_{r+2} w''\} \\ \cup \{v_j w' \mid r+3 \leq j \leq n-k\}$$

if $k \equiv 1 \pmod{4}$; and

$$E' = \{v_i w_{2i-1}, v_i w_{2i} \mid 1 \leq i \leq r\} \cup \{v_{r+1} w, v_{r+2} w''\} \\ \cup \{v_j w' \mid r+3 \leq j \leq n-k\}$$

if $k \equiv 3 \pmod{4}$.

In Fig. 2 we display $F_{14,10}$, $F_{15,11}$, $F_{16,12}$ and $F_{18,13}$.




Fig. 2. The graphs $F_{14,10}$, $F_{15,11}$, $F_{16,12}$ and $F_{18,13}$.

Proposition 2.4. *For any two positive integers n and k with $10 \leq k \leq \frac{4n}{5}$, there exists a graph G of order n with $\gamma_{t[1,2]}(G) = k$.*

Proof. Note that the order of the graph $F_{n,k}$ as constructed above is n and

$$W = \begin{cases} \{w_1, \dots, w_{\frac{k}{2}}\} \cup \{w'_1, \dots, w'_{\frac{k}{2}}\}, & k \equiv 0 \pmod{2} \\ \{w_1, \dots, w_{\frac{k-3}{2}}\} \cup \{w'_1, \dots, w'_{\frac{k-3}{2}}\} \cup \{w, w', w''\}, & k \equiv 1 \pmod{2}, \end{cases}$$

is a total $[1, 2]$ -set of G . So, $\gamma_{t[1,2]}(F_{n,k}) \leq k$. Take a $\gamma_{t[1,2]}$ -set S of $F_{n,k}$.

Case 1. $k \equiv 0 \pmod{2}$.

Since S is a total dominating set, $\{w_1, w_2, \dots, w_{\frac{k-2}{2}}\} \subseteq S$. On the other hand, $V(K_{n-k}) \cap S = \emptyset$. Otherwise, if $v_i \in S$ for some $1 \leq i \leq n-k$, then $\{v_i, w_1, w_2\} \subseteq N(v_1) \cap S$ if $i \neq 1$ or $\{v_i, w_3, w_4\} \subseteq N(v_2) \cap S$ if $i = 1$, both contradicting that S is a total $[1, 2]$ -set of G . Then Combining the above two facts with the other fact that $\{w_1, w_2, \dots, w_{\frac{k}{2}}\}$ is an independent set of $F_{n,k}$, it follows that $S = \{w_1, w_2, \dots, w_{\frac{k}{2}}\} \cup \{w'_1, w'_2, \dots, w'_{\frac{k}{2}}\}$, and thus $\gamma_{t[1,2]}(F_{n,k}) = k$.

Case 2. $k \equiv 1 \pmod{2}$.

Since S is a total dominating set, $\{w_1, w_2, \dots, w_{\frac{k-3}{2}}\} \subseteq S$. On the other hand, $V(K_{n-k}) \cap S = \emptyset$. Otherwise, if $v_i \in S$ for some $1 \leq i \leq n-k$, then $\{v_i, w_1, w_2\} \subseteq N(v_1) \cap S$ if $i \neq 1$ or $\{v_i, w_3, w_4\} \subseteq N(v_2) \cap S$ if $i = 1$, both contradicting that S is a total $[1, 2]$ -set. Then Combining the above two facts with the other fact that $\{w_1, w_2, \dots, w_{\frac{k-3}{2}}, w, w''\}$ is an independent set of $F_{n,k}$, it follows that $S = \{w_1, w_2, \dots, w_{\frac{k-3}{2}}\} \cup \{w'_1, w'_2, \dots, w'_{\frac{k-3}{2}}\} \cup \{w, w', w''\}$, and thus $\gamma_{t[1,2]}(F_{n,k}) = k$. $\square$

The upper bound on $\gamma_{t[1,2]}(G)$ in Theorem 2.3 is sharp. However, every extremal graph G with $\gamma_{t[1,2]}(G) = \frac{4n}{5}$ satisfies that $\delta(G) = 1$. It is natural to consider the upper bound on $\gamma_{t[1,2]}(G)$ when $\delta(G) \geq 2$.

Theorem 2.5. *Let G be a connected graph of order $n \geq 3$ and $\delta(G) \geq 2$. If $\gamma_{t[1,2]}(G) < +\infty$, then*

$$\gamma_{t[1,2]}(G) \leq \frac{2n}{3}.$$

Proof. Let S be a $\gamma_{t[1,2]}(G)$ -set of G . Same as the proof of Theorem 2.3, we divide S into four subsets S_1, S_2, S_3 and S_4 .

Claim 1. (1) $|pri(S_1, S)| \geq |S_1|$, (2) $|pri(S_3, S)| \geq \frac{2|S_3|}{3}$, (3) $|pri(S_4, S)| \geq |S_4| - 2\omega_4$, where ω_4 denotes the number of components in $G[S]$, which is isomorphic to a path of order at least four.

Claim 2. $|U| \geq \frac{|S_2|}{2}$ where $U = V(G) \setminus (S \cup pri(S_1, S) \cup pri(S_3, S) \cup pri(S_4, S))$.

Proof of Claim 2. Since $\delta(G) \geq 2$, every vertex in S_2 has at least one neighbor in U . Let $W = \cup_{u \in S_2} N(u) \cap U$. Since every vertex in W is adjacent

to at most 2 vertices in S and $\delta(G) \geq 2$, we have

$$2|W| \geq |E[S, W]| \geq |E[S_2, W]| \geq \sum_{v \in S_2} (d(v) - 1) \geq \sum_{v \in S_2} 1 \geq |S_2|,$$

and thus $|U| \geq |W| \geq \frac{|S_2|}{2}$. □

Since $\omega_4 \leq \frac{|S_4|}{4}$, we have

$$\begin{aligned} n &= |S| + |pri(S_1, S)| + |pri(S_3, S)| + |pri(S_4, S)| + |U| \\ &\geq |S| + |S_1| + \frac{2|S_3|}{3} + |S_4| - 2\omega_4 + \frac{|S_2|}{2} \\ &\geq \frac{3|S|}{2} + \frac{|S_1|}{2} + \frac{|S_3|}{6} \\ &\geq \frac{3|S|}{2}. \end{aligned}$$

So $\gamma_{t[1,2]}(G) \leq \frac{2n}{3}$ and the equality holds only if $S_1 = S_3 = \emptyset$, $S_4 = 4\omega_4$, $|pri(S_4)| = \frac{|S_4|}{2}$ and $|U| = |W| = \frac{|S_2|}{2}$. □

This upper bound is also sharp. For an integer $k \geq 4$, we can construct a graph F_k of order $n = 3k$ with $\gamma_{t[1,2]}(F_k) = 2k$ as follows. We start from the complete graph K_k with $V(K_k) = \{v_1, v_2, \dots, v_k\}$. Let $V(F_k) = V(K_k) \cup W \cup W'$, where $W = \{w_1, w_2, \dots, w_k\}$ and $W' = \{w'_1, w'_2, \dots, w'_k\}$, and $E(F_k) = \cup_{i=1}^k \{v_i w_i, w_i w'_i, w'_i v_{i+1}\} \cup E(K_k)$, where $i + 1$ is taken modulo k . Therefore, F_k is a graph of order $3k$ and for any $1 \leq i \leq k$, $d(v_i) = k + 1$ and $d(w_i) = d(w'_i) = 2$. The graph F_4 is shown in Fig. 3.




Fig. 3. The graph F_4

Proposition 2.6. For an integer $k \geq 4$, $\gamma_{t[1,2]}(F_k) = \frac{2n}{3}$, where $n = 3k$.

Proof. Let $S = W \cup W'$. One can see that S is a total $[1, 2]$ -set of F_k and thus $\gamma_{t[1,2]}(F_k) \leq \frac{2n}{3}$. Now suppose that $\gamma_{t[1,2]}(F_k) < \frac{2n}{3}$ and S' is a $\gamma_{t[1,2]}$ -set of F_k . Then $S' \cap V(K_k) \neq \emptyset$ and $|S' \cap V(K_k)| \leq 2$.

Assume first that $|V(K_k) \cap S'| = 2$ and let $v_i, v_j \in S'$ for some two distinct $i, j \in \{1, 2, \dots, k\}$. In this case, $(\{v_1, v_2, \dots, v_k\} \setminus \{v_i, v_j\}) \cap S' = \emptyset$ and thus $w_l, w'_l \in S'$ for any $l \in \{1, 2, \dots, k\} \setminus \{i, j\}$. But, $|N_G(v_l) \cap S'| \geq 3$, a contradiction. Now assume that $|V(K_k) \cap S'| = 1$ and $V(K_k) \cap S' = \{v_1\}$. Since $v_i \notin S'$ for any $i \in \{2, 3, 4\}$, we have $w_2, w'_2 \in S'$ and $w_3, w'_3 \in S'$, and thus $|N_G(v_3) \cap S'| \geq 3$, a contradiction. $\square$

Now, for $8 \leq k \leq \frac{2n}{3} - 1$, we construct a graph $H_{n,k}$ of order n , $\gamma_{t[1,2]}(H_{n,k}) = k$ and $\delta(H_{n,k}) \geq 2$ as follows. We start from the complete graph K_{n-k} with $V(K_{n-k}) = \{v_1, v_2, \dots, v_{n-k}\}$ and $r = \lfloor \frac{k-2}{2} \rfloor$. The graph $H_{n,k}$ has $V(H_{n,k}) = V(K_{n-k}) \cup \{w_1, \dots, w_r\} \cup \{w'_1, \dots, w'_r\} \cup V'$, where

$$V' = \begin{cases} \{w, w'\}, & \text{if } k = 2r + 2 \\ \{w, w', w''\}, & \text{if } k = 2r + 3, \end{cases}$$

and

$$E(H_{n,k}) = E(K_{n-k}) \cup \{v_i w_i, w_i w'_i, w'_i v_{i+1} \mid 1 \leq i \leq r-1\} \cup \{v_r w_r, w_r w'_r, w'_r v_1\} \cup E',$$

where

$$E' = \{w w'\} \cup \{v_j w, v_j w' \mid r+1 \leq j \leq n-k\},$$

if $k = 2r + 2$; and

$$E' = \{v_{r+1} w, v_{r+2} w'', w w', w' w''\} \cup \{v_j w, v_j w'' \mid r+3 \leq j \leq n-k\},$$

if $k = 2r + 3$. See Fig. 4 for an illustration for $H_{14,8}$ and $H_{15,9}$.




Fig. 4. The graphs $H_{14,8}$ and $H_{15,9}$

Proposition 2.7. *For any n and k with $8 \leq k \leq \frac{2n}{3} - 1$, there exists a graph G of order n with $\delta(G) \geq 2$ and $\gamma_{t[1,2]}(G) = k$.*

Proof. Note that the order of the graph $H_{n,k}$ as constructed above is n and

$$S' = \begin{cases} \{w_1, \dots, w_r\} \cup \{w'_1, \dots, w'_r\} \cup \{w, w'\}, & \text{if } k = 2r + 2 \\ \{w_1, \dots, w_r\} \cup \{w'_1, \dots, w'_r\} \cup \{w, w', w''\}, & \text{if } k = 2r + 3, \end{cases}$$

is a total $[1, 2]$ -set of $H_{n,k}$. So, $\gamma_{t[1,2]}(H_{n,k}) \leq k$. In addition, any proper subset of S' is not a total $[1, 2]$ -set of $H_{n,k}$. Let S be a $\gamma_{t[1,2]}$ -set of $H_{n,k}$. Note first that $|S \cap V(K_{n-k})| \leq 2$ since S is a $[1, 2]$ -set of $H_{n,k}$. Our proof is based on the fact that: if $v_i \notin S$ and $v_{i+1} \notin S$ for $1 \leq i \leq r - 1$, then $\{w_i, w'_i\} \subseteq S$ since $N(w_i) = \{v_i, w'_i\}$, $N(w'_i) = \{w_i, v_{i+1}\}$ and S is a total $[1, 2]$ -set; and similarly if $S \cap \{v_{r+1}, v_{r+2}, \dots, v_{n-k}\} = \emptyset$, then $\{w, w'\} \subseteq S$ if $k = 2r + 2$ or $\{w, w', w''\} \subseteq S$ if $k = 2r + 3$.

Claim 1. $|S \cap V(K_{n-k})| \leq 1$.

By contradiction, suppose that $|S \cap V(K_{n-k})| = 2$ and $v_l, v_j \in S$. If $|S \cap \{v_1, \dots, v_r\}| \leq 1$, there exists $1 \leq i \leq r$ such that $v_i, v_{i+1} \notin S$ since $r \geq 3$, then $\{w_i, w'_i\} \subseteq S$. But now, $\{w_i, v_l, v_j\} \subseteq N(v_i) \cap S$. This contradicts to the fact that S is a $[1, 2]$ -set. So we may assume $|S \cap \{v_1, \dots, v_r\}| = 2$. Then $w \in S$ since $\{v_i \mid r+1 \leq i \leq n-k\} \cap S = \emptyset$, which means $\{w, v_l, v_j\} \subseteq N(v_{r+1}) \cap S$, a contradiction.

Claim 2. $S \cap V(K_{n-k}) = \emptyset$.

By contradiction, suppose that $|S \cap V(K_{n-k})| = 1$ and $v_l \in S \cap V(K_{n-k})$. If $1 \leq l \leq r$, then $\{w, w'\} \subseteq S$, and then $\{w, w', v_l\} \subseteq N(v_{r+1}) \cap S$ if $k = 2r + 2$ or $\{w, w'', v_l\} \subseteq N(v_{r+3}) \cap S$ if $k = 2r + 3$, both contradicting to the fact that S is a $[1, 2]$ -set. If $r+1 \leq l \leq n-k$, then $\{w_1, w'_1, w_2, w'_2\} \subseteq S$, so $\{w'_1, w_2, v_l\} \subseteq N(v_2) \cap S$, a contradiction.

It follows immediately that $S = S'$ and thus $\gamma_{t[1,2]}(H_{n,k}) = k$. $\square$

3 Graphs with no total $[1, 2]$ -set

As we have seen in the previous section, there exist many graphs with no total $[1, 2]$ -set, we summarize it as follows.

Theorem 3.1. *Let G be a connected graph of order n .*

- (1) *If $n \geq 3$ and $\gamma_{[1,2]}(G) > \frac{4n}{5}$, then $\gamma_{t[1,2]}(G) = +\infty$;*
- (2) *If $n \geq 5$, $\delta(G) \geq 2$ and $\gamma_{[1,2]}(G) > \frac{2n}{3}$, then $\gamma_{t[1,2]}(G) = +\infty$;*
- (3) *Let G be a tree of order n with k leaves. If $\gamma_{[1,2]}(G) = n - k$, then $\gamma_{t[1,2]}(G) = +\infty$ unless G is a caterpillar.*

Proof. (1) and (2) are immediate from Theorem 2.3 and 2.6.

(3) Assume that $\gamma_{t[1,2]}(T)$ exists and S is a $\gamma_{t[1,2]}(T)$ -set of T . Let L be the set of all leaves of T . If $v \in L \cap S$, the support vertex of v also lies in S . It follows that $S' = S \setminus L$ is also a $[1, 2]$ -set of T and hence $|S'| \leq n - k$. Since $\gamma_{[1,2]}(T) = n - k$, we have $|S'| = n - k$ and $V(T) \setminus S' = L$. If there exists a vertex $v \in S'$ such that $N(v) \subseteq S'$, then $S' \setminus \{v\}$ is a $[1, 2]$ -set of T with cardinality less than $n - k$, a contradiction. So each vertex in S' must have at least one neighbor outside S' , which means all the vertices of S' are support vertices of T . It is easy to see $\Delta(T[S']) \leq \Delta(T[S]) \leq 2$. So $T[S']$ must be a path and then T is a caterpillar. $\square$

Let p and k be two integers with $p \geq k + 2 \geq 5$, $G_{p,k}$ is a graph obtained from a complete graph K_p as follows: for every k -element subsets S of the vertices set $V(K_p)$, we add a new vertex x_s and the edges $x_s u$ for all $u \in S$. In [14], Yang and Wu proved that $\gamma_{[1,2]}(G_{p,k}) = |V(G_{p,k})|$. So, it is immediate from Theorem 3.1 that

Corollary 3.2. *If $p \geq k + 2$ and $k \geq 3$, then $\gamma_{t[1,2]}(G_{p,k}) = +\infty$.*

4 Graphs with $\gamma_{t[1,2]}(G) = \gamma_{[1,2]}(G)$

It was shown in [4] that if G is the corona $H \circ K_1$ of a graph H , then $\gamma_{[1,2]}(G) = \gamma(G)$. For total $[1, 2]$ -domination number, we proved the following result.

Theorem 4.1. *Let G be the corona $H \circ K_1$ of a connected graph H of order $n \geq 2$. Then $\gamma_{t[1,2]}(G) = \gamma_{[1,2]}(G)$ if and only if H is a path or a cycle.*

Proof. To show the sufficiency, let S be a $\gamma_{t[1,2]}(G)$ -set of G . Since S is a dominating set, for a leaf of G , either itself is contained in S or its support vertex contained in S . Moreover, By the definition of corona, for two leaves, their support vertices are different. So, $|S| \geq n$. On the other hand, since H is a path or cycle, $V(H)$ is a total $[1, 2]$ -set of G , $|S| \leq |V(H)| = n$. This proves the sufficiency.

To show the necessity, we assume that $\gamma_{t[1,2]}(H \circ K_1) = \gamma_{[1,2]}(H \circ K_1)$, and let S be a $\gamma_{t[1,2]}(G)$ -set of G . Obviously, all the support vertices must lie in S , which means $V(H) \subseteq S$. On the other hand, $\Delta(G[S]) \leq 2$, implying that $\Delta(H) \leq 2$. Moreover, since H is connected, $H \cong P_n$ or $H \cong C_n$. $\square$

Theorem 4.2. ([2]) *If G is a graph without isolated vertices, then G has a minimum dominating set D such that, for all $v \in D$, $\text{pri}(v, D) \cap (V(G) \setminus D) \neq \emptyset$.*

Using the above result, Chellali et al. [4] gave the following sufficient condition for a graph G satisfying $\gamma(G) = \gamma_{[1,2]}(G)$.

Theorem 4.3. ([4]) *If G is a P_4 -free graph, then $\gamma(G) = \gamma_{[1,2]}(G)$.*

By slightly simplifying the proof of the above theorem, we get the following stronger result.

Theorem 4.4. *Let G be a P_4 -free graph without isolated vertices. If D is a minimum dominating set of G such that $\text{pri}(v, D) \cap (V(G) \setminus D) \neq \emptyset$ for all $v \in D$, then D is also a $[1, 2]$ -set of G .*

Proof. By contradiction, suppose that D is not a $[1, 2]$ -set of G . Then there exists a vertex $u \in V \setminus D$ with at least three neighbors x, y, z say, in D . For convenience, we use A_v to denote the set of private neighbors of v in $V \setminus D$ for a vertex $v \in D$. Hence, $A_v \neq \emptyset$ for every $v \in \{x, y, z\}$. We consider two cases.

Case 1. $A_x \cup A_y \cup A_z \not\subseteq N(u)$.

Without loss of generality, let $z' \in A_z \setminus N(u)$. If $xz \notin E(G)$, then $\{z', z, u, x\}$ induces a P_4 in G . So, $xz \in E(G)$. Similarly, we have $yz \in E(G)$. Furthermore, $xy \in E(G)$, since otherwise, $\{x', x, z, y\}$ induces a P_4 in G , where $x' \in A_x$. To avoid a P_4 induced by $\{z', z, x, x'\}$ in G , we have $x'z' \in E(G)$. But, $\{y, x, x', z'\}$ will induce a P_4 in G , a contradiction.

Case 2. $A_x \cup A_y \cup A_z \subseteq N(z)$.

Note that $(N(y) \cap N(z)) \setminus (N(u) \cup N(x)) \neq \emptyset$. If this is not, then $(D \setminus \{y, z\}) \cup \{u\}$ is a dominating set of G with cardinality less than $|D|$, a contradiction. By a similar argument to the above, we have $(N(x) \cap N(y)) \setminus (N(u) \cup N(z)) \neq \emptyset$. Take $p \in (N(y) \cap N(z)) \setminus (N(u) \cup N(x))$ and $q \in (N(x) \cap N(y)) \setminus (N(u) \cup N(z))$ respectively.

If $pq \in E(G)$, then each of $\{u, z, p, q\}$ induces P_4 in G . So, $pq \notin E(G)$. If $xy \notin E(G)$, then $pyqx$ induces a P_4 ; if $yz \notin E(G)$, then $qypz$ induces a P_4 in G . So, it follows that $xy \in E(G)$ and $yz \in E(G)$. By a similar argument to the above, one has $xz \in E(G)$. But, then $\{q, x, z, p\}$ induces P_4 , a contradiction.

Consequently, D is a $[1, 2]$ -set and the theorem is proved. $\square$

Theorem 4.5. *If G is a connected P_4 -free graph of order $n \geq 4$, then*

$$\gamma_{[1,2]}(G) = \begin{cases} 2, & \text{if } \Delta(G) = n - 1 \\ \gamma_{[1,2]}(G) = \gamma(G), & \text{if } \Delta(G) < n - 1. \end{cases}$$

Proof. If $\Delta(G) = n - 1$, then the result trivially holds. So, next we assume that $\Delta(G) < n - 1$. By Theorem 4.4, let D be a $\gamma_{[1,2]}(G)$ -set of G such that $\text{pri}(u, D) \cap (V(G) \setminus D) \neq \emptyset$ for all $u \in D$. Among all such dominating sets, we choose one such that $\omega(G[D])$ is as small as possible. We show that D is also a $\gamma_{t[1,2]}(G)$ -set of G . By contradiction, suppose that D is not a total $[1, 2]$ -set. Then, either $\delta(G[D]) = 0$ or $\Delta(G[D]) \geq 3$.

Case 1. $\Delta([D]) \geq 3$.

Suppose $v \in D$ is a vertex having three neighbors v_1, v_2, v_3 say, in D , and let $w \in \text{pri}(v, D)$. $w_i \in \text{pri}(v_i, D)$ for $i = 1, 2, 3$. To avoid having a P_4 induced by $\{w_1, v_1, v, v_2\}$, we have $v_1 v_2 \in E(G)$. Similarly, $ww_2 \in E(G)$, since otherwise $\{w, v, v_2, w_2\}$ induces a P_4 . But then $\{w, w_2, v_2, v_1\}$ induces a P_4 , a contradiction.

Case 2. $\Delta(G[D]) \leq 2$ and $\delta(G[D]) = 0$.

Since $\Delta([D]) \leq 2$, all components of $G[D]$ are paths or cycles. Combing this with the fact that G is P_4 -free, each component of $G[D]$ is isomorphic to K_1, K_2, P_3, K_3 or C_4 . Without loss of generality, let H_i be an isolated vertex for any integer $i \leq k$ and H_j is a path of order at least two or a cycle for any $k + 1 \leq j \leq t$. Since G is P_4 -free, $\text{diam}(G) \leq 2$, which implies that $N(x) \cap N(y) \neq \emptyset$ if x and y lie two different components of $G[D]$. Without loss of generality, let $V(H_1) = \{v_1\}$. Let $v_1 w v_j$ be a path connecting v_1 and a vertex $v_j \in V(H_j)$ for some $j \geq 2$. To avoid having a P_4 induced by $\{v_1, w, v_j, v'_j\}$ for any $v'_j \in N(v_j) \cap V(H_j)$, and thus $V(H_j) \subseteq N(v_1)$. But then $D' = (D \setminus V(H_j)) \cup \{w\}$ is a dominating set of G with cardinality less than $|D|$ if $|V(H_j)| \geq 2$. So, $|V(H_j)| = 1$ for any j , i.e. each component of $G[D]$ is an isolated vertex.

Since $\Delta(G) < n - 1$, $\gamma_{t[1,2]}(G) = k \geq 2$. Let $D = \{v_1, v_2, \dots, v_k\}$. Since G is connected, $N(v_1) \cap N(v_j) \neq \emptyset$ for some integer $j \in \{2, \dots, k\}$. Without loss of generality, let $j = 2$. Take $w \in N(v_1) \cap N(v_2)$ and $w_i \in \text{pri}(v_i, D)$ for $i \in \{1, 2\}$. To avoid a P_4 , $\text{pri}(v_1, D) \cup \text{pri}(v_2, D) \subseteq N(w)$, since otherwise, if $w_1 w \notin E(G)$, $\{w_1, v_1, w, v_2\}$ induces a P_4 . If $(N(v_1) \cap N(v_2)) \setminus \{w\} \subseteq N(w)$, then $(D \setminus \{v_1, v_2\}) \cup \{w\}$ is a dominating set of G , a contradiction. If $(N(v_1) \cap N(v_2)) \setminus \{w\} \not\subseteq N(w)$, then $D' = (D \setminus \{v_2\}) \cup \{w\}$ is a dominating set of G . Since $(N(v_1) \cap N(v_2)) \setminus \{w\} \not\subseteq N(w)$, $\text{pri}(v_1, D') \neq \emptyset$; $v_2 \in \text{pri}(w, D')$, $\text{pri}(v_i, D') = \text{pri}(v_i, D)$ for $3 \leq i \leq k$ and $\omega(G[D']) < \omega(G[D])$, contradicting the choice of D that $\omega(G[D])$ is as small as possible.

□

5 Further research

It is interesting that the total $[1, 2]$ -domination problem is concerned with graph partition and factors. Recall that a spanning subgraph H of G is called a $[a, b]$ -factor if $a \leq d_H(v) \leq b$. In particular, H is called a k -factor of G if H is a k -regular spanning subgraph of G .

Conjecture 1. For any cubic graph G of order n , $\gamma_{t[1,2]}(G) < n$.

The statement of Conjecture 1 is equivalent to that every cubic graph G has a vertex partition $(S, V(G) \setminus S)$ such that $1 \leq \delta(G[S]) \leq \Delta(G[S]) \leq 2$ and $1 \leq \delta(G - S) \leq \Delta(G - S) \leq 2$.

It is well-known that every regular graph has a $[1, 2]$ -factor (see [11]), and so does a cubic graph. Hence Conjecture 1 asserts the existence of $[1, 2]$ -factor with an additional property in a cubic graph. A theorem of Petersen [10] says that every even regular graph G has a 2-factor. So, we also pose the following conjecture.

Conjecture 2. For any 4-regular graph G of order n , $\gamma_{t[1,2]}(G) < n$.

The middle levels problem, attributed to Håvel [8], concerns the following family of graphs. Let n, a, b be integers with $0 \leq a < b \leq n$. Let $G(n; a, b)$ denote the bipartite graph whose vertices are all the a -element and b -element subsets of an n -set, say $[n] = \{1, 2, \dots, n\}$. An a -element subset A and a b -element subset B are adjacent in $G(n; a, b)$ if and only if $A \subseteq B$. So, the order of $G(n; a, b)$ is $\binom{n}{a} + \binom{n}{b}$. The middle levels problem asks that for a positive integer k , is the graph $G(2k+1; k, k+1)$ Hamiltonian? It is so-named because it deals with the central levels of the Boolean algebra $2^{[2k+1]}$. Yang and Wu [14] proved the following theorem.

Theorem 5.1. ([14]) *Let n and k be two integers with $n \geq k \geq 3$. If n is sufficiently large with respect to any fixed k , then $\gamma_{[1,2]}(G(n; k, n-k)) = |V(G(n; k, n-k))|$.*

So, by Theorems 3.1 and 5.1, if n is sufficiently large with respect to any fixed k , then $\gamma_{t[1,2]}(G(n; k, n-k)) = +\infty$. Note that $G(n; k, n-k)$ is a $\binom{n-k}{k}$ -regular bipartite graph of order $2\binom{n}{k}$. So, we ask the following problem.

Question. What is the smallest integer k such that there exists a k -regular graph G with $\gamma_{t[1,2]}(G) = +\infty$?

Chellali et al. [4] asked that if $\gamma_{[1,2]}(G) < n$ for any graph 5-regular graph G ? If this is true, it is equivalent to the following conjecture.

Conjecture 3. Every 5-regular graph G has an induced subgraph H with $3 \leq d_H(v) \leq 4$.

Note that a stronger assertion, that every 5-regular graph G has an 3-regular or 4-regular induced subgraph H , generally does not hold. The formulation of Conjecture 3 reminds us the well-known Berge-Saure conjecture [1], confirmed by Tâskinov [12] and Zhang [15], which says that every 4-regular graph contains a 3-regular subgraph.

References

- [1] N. Alon, S. Friedland, and G. Kalai, Regular subgraphs of almost regular graphs, *J. Combin. Theory Ser. B* 37(1984), 79-91.
- [2] B. Bollobás and E.J. Cockayne, Graph-theoretic parameters concerning domination, independence, and irredundance, *J. Graph Theory* 3(1979), 241-249.
- [3] R.C. Brigham, J.R. Carrington and R.P. Vitray, Connected graphs with maximum total domination number, *J. Combin. Comput. Combin. Math.* 34(2000), 81-96.
- [4] M. Chellali, T.W. Haynes and S.T. Hedetniemi, $[1, 2]$ -set in graphs, *Discrete Appl. Math.* 161(2013), 2885-2893.
- [5] E.J. Cockayne, R. Dawes and S.T. Hedetniemi, Total domination in graphs, *Networks* 10(1980), 211-215.
- [6] I.J. Dejter, Quasiperfect domination in triangular lattices, *Discuss. Math. Graph Theory* 29(2009), 179-198.
- [7] M. Fischermann, Unique total domination graphs, *Ars Combin.* 73(2004), 289-297.
- [8] I. Håvel, Semipaths in directed cubes, in: M. Fiedler (Ed.), *Graphs and Other Combinatorial Topics*, Teubner, Leipzig, 1983, pp. 101-108.
- [9] M.A. Henning, A survey of selected recent results on total domination in graphs, *Discrete Math.* 309(1009), 32-63.
- [10] J. Petersen, Die Theorie der regulären Graph, *Acta Math.* 15(1981), 193-220.
- [11] E.R. Sheinerman and D.H. Ullman, *Fractional Graph Theory*, Wiley, New York (1997).

- [12] V.A. Tâskinov, Regular subgraphs of regular graphs, *Soviet Math. Dokl.* 26(1982), 37-38.
- [13] D.B. West, Introduction to Graph Theory, Second Edition, Prentice Hall, Upper Saddle River, NJ(2001).
- [14] X. Yang and B. Wu, $[1, 2]$ -domination in graphs, *Discrete Appl. Math.* 175(2014), 79-86.
- [15] L. Zhang, Every 4-regular simple graph contains a 3-regular subgraph, *J. Changsha Railway Inst.* 1(1985) 130-154.